

AD-A166 606

ONE-COMPONENT PLASMA MODEL AND LARGE Z LABORATORY  
PLASMAS(U) BATTELLE COLUMBUS LABS ON R L LIBOFF ET AL.  
APR 86 ARFSD-CR-86005 DARG29-81-D-0100


1/1

UNCLASSIFIED

F/G 20/9

NL





MICROCOPY

CHART

12

AD-A166 606

AD

~~AD-E401-408~~

CONTRACTOR REPORT ARFSD-CR-86005

**ONE-COMPONENT PLASMA MODEL AND  
LARGE Z LABORATORY PLASMAS**

**RICHARD L. LIBOFF**  
SCHOOL OF ELECTRICAL ENGINEERING  
AND APPLIED PHYSICS  
CORNELL UNIVERSITY  
ITHACA, NY 14853

**G. L. FERRENTINO**  
PROJECT ENGINEER

**DTIC**  
**ELECTE**  
APR 14 1986  
**S** **D**  
**E**

**APRIL 1986**

  
US ARMY  
ARMAMENT  
MUNITIONS &  
CHEMICAL COMMAND  
ARMAMENT R&D CENTER

**U. S. ARMY ARMAMENT RESEARCH AND DEVELOPMENT CENTER**

**FIRE SUPPORT ARMAMENT CENTER**

**DOVER, NEW JERSEY**

**APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.**

**DTIC FILE COPY**

The views, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

Destroy this report when no longer needed. Do not return to the originator.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
Contractor Report ARFSD-CR-86005	AD-A166606	
4. TITLE (and Subtitle)		5. TYPE OF REPORT & PERIOD COVERED
ONE-COMPONENT PLASMA MODEL AND LARGE Z LABORATORY PLASMAS		Interim Aug 85 to Dec 85
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)		8. CONTRACT OR GRANT NUMBER(s)
Richard L. Liboff, Cornell University G. L. Ferrentino, Project Engineer, ARDC		DAA G29-81-D-0100 (D.O. No. 1769, TCN 85-526)
9. PERFORMING ORGANIZATION NAME AND ADDRESS		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
School of Electrical Engineering and Applied Physics Cornell University Ithaca, NY 14853		
11. CONTROLLING OFFICE NAME AND ADDRESS		12. REPORT DATE
ARDC, IMD STINFO Div (SMCAR-MSI) Dover, NJ 07801-5001 (cont)		April 1986
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES
ARDC, FSAC Artillery Armament Div (SMCAR-FSA-E) Dover, NJ 07801-5001		
		15. SECURITY CLASS. (of this report)
		Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)		
Approved for public release; distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
This research was supported in part by Battelle Columbus Laboratory.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Railgun, Thermodynamic Plasma, Internal energy Pressure, Recombination Temperature, Electrons Equation of state		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
Stemming from an expression for the mean two-particle potential energy of a two-component laboratory plasma comprised of ions of charge Ze and electrons, it is argued that for sufficiently large Z, thermodynamic properties of such plasmas are the same as those relevant to Wigner's one-component plasma model. Thus, an equation of state is obtained for a strongly coupled laboratory plasma with $Z \geq 5$ .		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

11. CONTROLLING OFFICE NAME AND ADDRESS: (cont)

Commander

U.S. Army Research Office

ATTN: AMCRO-PR

P.O. Box 12211

Research Triangle Park, NC 27709

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

## CONTENTS

	Page
Introduction	1
Discussion	2
References	9
Distribution List	11

# INTRODUCTION

Stemming from an expression for the mean two-particle potential energy of a two-component laboratory plasma comprised of ions of charge  $Ze$  and electrons, it is argued that for sufficiently large  $Z$ , thermodynamic properties of such plasmas are the same as those relevant to Wigner's one-component plasma model. Thus the following equation of state is obtained for a laboratory plasma with  $Z \geq 5$  and  $\Gamma \geq 1$ .

$$\frac{P}{nk_B T} = 1 + \frac{d}{3} + \frac{1}{3} (\bar{a}\Gamma + b\Gamma^{1/4} + c\Gamma^{-1/4})$$

In this equation,  $P$  is pressure,  $T$  is temperature,  $n$  is ion number density and  $\bar{a}$ ,  $b$ ,  $c$ ,  $d$  are known numerical constants. The plasma parameter  $\Gamma = (Ze)^2 / ak_B T$  where  $4\pi a^3 / 3$  represents mean occupation volume per ion.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	





## DISCUSSION

Strongly coupled plasmas play a role in recombination approaches to x-ray lasing [1], inertial-confinement fusion devices [2], the interiors of certain super-dense stars [3], and in plasma-driven rail-gun devices [4].

For the most part studies addressing strongly coupled plasmas [5,6,7] have employed a plasma model due to Wigner [8] which was conceived for the purposes of studying phase transition to the solid state. This medium is called "jellium", or more commonly, a "one-component plasma", which often carries the abbreviation, OCP. An OCP is comprised of ions moving in a charge-neutralizing uniform negative background.

In the present work attention is directed at a laboratory plasma comprised of ions of charge  $Ze$  and electrons in equilibrium at a given temperature. Examining the interaction energy of the plasma indicates that for  $Z \geq 5$ , thermodynamic properties such as internal energy and equation of state are given by corresponding expressions appropriate to an OCP. As an application of this finding, the equation of state for such a relatively high  $Z$  plasma is obtained from a previously constructed expression for the Helmholtz free energy [9].

With the plasma under consideration comprised of electrons and ions of charge  $eZ$ , charge neutrality implies the constraint

$$eV(n_e - Zn_Z) \approx 0 \quad (1)$$

where  $V$  is plasma volume and  $n_e$  and  $n_Z$  are electron and ion number densities, respectively. The approximate equality in

(1) derives from the inherent statistical nature of a plasma. For a two-component plasma, three interactions contribute to the mean two-particle potential energy and we write

$$\langle V \rangle \approx (4\pi)^{1/3} [n_e^{1/3} e^2 - (n_e n_Z)^{1/6} Z e^2 + n_Z^{1/3} (Ze)^2] \quad (2)$$

With the constraint (1) we may set  $n_e = Z n_Z$  and (2) becomes

$$\langle V \rangle \approx e^2 (4\pi n_Z)^{1/3} (Z^{1/3} - Z^{7/6} + Z^2) \quad (3)$$

Thus in the limit

$$Z^2 \gg Z^{1.17} \quad (4)$$

the relation (3) reduces to

$$\langle V \rangle \approx (4\pi n_Z)^{1/3} (Ze)^2 \quad (5)$$

which we recognize to be the interaction potential of a one-component plasma comprised of ions of charge  $Ze$ .

This similarity may be further illustrated through the plasma parameter [10,11]

$$\gamma^{2/3} \approx \frac{\langle V \rangle}{\langle E_K \rangle} \approx \frac{\langle V \rangle}{k_B T} \quad (6)$$

where  $\langle E_K \rangle$  is mean kinetic energy per particle. The relevance of  $\gamma$  to the properties of a plasma is evident from (6). Namely, with this expression we may conclude that a plasma is strongly coupled when  $\gamma \gtrsim 1$  and weakly coupled when  $\gamma \ll 1$ .

Substituting (5) into (6) gives (dropping the  $Z$  subscript on  $n_Z$ )

$$\gamma^{2/3} \approx \frac{(Ze)^2 (4\pi n)^{1/3}}{k_B T} \quad (7)$$

which, again, is the plasma parameter relevant to an OCP with ions of charge  $Ze$ .

The canonical expression for  $\gamma$  is given by

$$\gamma = \frac{1}{4\pi n \lambda_D^3} \quad (8)$$

where  $\lambda_D$  is the Debye distance.<sup>13,14</sup> With the latter two expressions we find

$$\lambda_D^2 = \frac{k_B T}{4\pi n (Ze)^2} \quad (9)$$

which is seen to be the Debye distance for an OCP comprised of ions of charge  $Ze$ .

These relations may be cast in terms of a plasma parameter more common to studies of OCP. It is given by

$$\Gamma = (Ze)^2 / a k_B T \quad (10)$$

where  $a^3$  is a measure of the mean occupation volume per ion.

That is,

$$\frac{4}{3} \pi a^3 n = 1 \quad (11)$$

The parallel structure of  $\gamma$  and  $\Gamma$  is evidenced by rewriting (8):

$$\gamma = \frac{(Ze)^2/\lambda_D}{k_B T} \quad (12a)$$

$$\Gamma = \frac{(Ze)^2/a}{k_B T} \quad (12b)$$

so that

$$\gamma^2 = 3\Gamma^3$$

We may conclude that the previously stated criterion that separates weakly from strongly coupled plasmas may also be given in terms of  $\Gamma$ .

Thus, we find that in the limit (4), the coupling and parameters of a two-component plasma of electrons and ions of charge  $Ze$  reduce to those relevant to an OCP of ions of charge  $Ze$  in a negative background. We may conclude that thermodynamic properties such as internal energy and equation of state for a two-component laboratory plasma with  $Z \gg Z^{1.17}$  are the same as those of an OCP comprised of the same species of ions.

Numerical work of Slattery, Doolan and DeWitt [9] established the following expression for internal energy  $U$  for the fluid phase of an OCP.

$$\frac{U}{Nk_B T} = \bar{a}\Gamma + b\Gamma^{1/4} + c\Gamma^{-1/4} + d + \bar{e}\Gamma/N \quad (13)$$

where  $N$  is total ion number and

$$\begin{aligned} \bar{a} &= -0.898, & b &= 0.950, & c &= 0.190, \\ d &= -0.815, & \bar{e} &= 0.010 \end{aligned}$$

The Helmholtz free energy,  $F$ , may then be obtained through integration. Namely,

$$\frac{F(\Gamma)}{Nk_B T} = \int_{\Gamma_1}^{\Gamma} \left[ \frac{U(\Gamma')}{Nk_B T} + 3 \right] d\Gamma' + \frac{F(\Gamma_1)}{Nk_B T} \quad (14)$$

The normalized free energy  $F(\Gamma_1)/Nk_B T$ , with  $\Gamma_1 = 1$  was calculated employing various contributions over the unit interval. There results (in the limit of large  $N$ ) [9]

(15)

$$\frac{F(\Gamma)}{Nk_B T} = \bar{a}\Gamma + 4(b\Gamma^{1/4} - c\Gamma^{-1/4}) + (d+3) \ln \Gamma - (\bar{a} + 4b - 4c + 1.152)$$

With this value of free energy at hand an equation of state is obtained from the thermodynamic relation

$$P = - \left( \frac{\partial F}{\partial V} \right)_T \quad (16)$$

To perform this differentiation we first rewrite  $\Gamma$  (10,12b) in explicit form

$$\Gamma = \frac{(Ze)^2}{k_B T} \left( \frac{4\pi N}{3V} \right)^{1/3} \quad (17)$$

There results

$$\frac{\partial \Gamma}{\partial V} = - \frac{\Gamma}{3V} \quad (18)$$

Differentiating (15) we obtain

$$\frac{1}{Nk_B T} \frac{\partial F}{\partial V} = [\bar{a} + b\Gamma^{-3/4} + c\Gamma^{-5/4} + (d+3)\Gamma^{-1}] \frac{\partial \Gamma}{\partial V} \quad (19)$$

With (16) and (18) we then obtain

$$\frac{PV}{Nk_B T} = 1 + \frac{d}{3} + \frac{1}{3} (\bar{a}\Gamma + b\Gamma^{1/4} + c\Gamma^{-1/4}) \quad (20)$$

With the previously stated argument we may conclude that (13) and (20) are valid energy and equation-of-state formulas for a laboratory plasma comprised of ions and electrons in equilibrium at a given temperature and obeying the constraint (4). Note in particular that for  $Z=5$  (completely ionized boron)  $Z^{1.17}/Z^2 = 0.26$ . For  $Z=13$  (completely ionized aluminum) this ratio becomes 0.12. Thus the proposed equivalence should be valid for  $Z \geq 5$ . It is also important to note that this equivalence is not appropriate to processes where the dynamics of electrons come into play, such as conductivity [12].

#### REFERENCES

1. D. M. Heffernan and R. L. Liboff, J. Plasma Phys. 27, 473 (1982).
2. K. Brueckner and S. Jorna, Revs. Mod. Phys. 46, 325 (1975).
3. E. E. Salpeter and H. M. Van Horn, Astrophys. J. 155, 183 (1969).
4. A collection of papers on electromagnetic propulsion is given in IEEE Trans. on Magnetics, Mag-20, March (1984).
5. S. Ichimaru, Revs. Mod. Phys. 54, 1017 (1982).
6. M. Baus and J. P. Hansen, Phys. Rep. 59, 1 (1980).
7. G. Kalman and P. Carini, Strongly Coupled Plasmas (Plenum, New York, 1978).
8. E. P. Wigner, Trans. Faraday Soc. (London) 34, 678 (1938).
9. W. L. Slattey, G. D. Doolen and H. E. Dewitt, Phys. Rev. A21, 2087 (1980); Phys. Rev. A26, 2255 (1982).
10. R. L. Liboff, J. Appl. Phys. 56, 2530 (1984)
11. P. M. Platzman and P. A. Wolf, Waves and Interactions in Solid State Plasmas (Academic, New York, 1973).
12. L. Spitzer, Jr., Physics of Fully Ionized Gases, (Interscience, New York, 1956).

DISTRIBUTION LIST

Commander  
Armament Research and Development Center  
U.S. Army Armament, Munitions  
and Chemical Command  
ATTN: SMCAR-FSA-E (5)  
SMCAR-MSI (5)  
SMCAR-SF  
Dover, NJ 07801-5001

Commander  
U.S. Army Armament, Munitions  
and Chemical Command  
ATTN: AMSMC-QAR-R(D)  
Dover, NJ 07801-5001

Administrator  
Defense Technical Information Center  
ATTN: Accessions Division (12)  
Cameron Station  
Alexandria, VA 22304-6145

Director  
U.S. Army Materiel Systems  
Analysis Activity  
ATTN: AMXSY-MP  
Aberdeen Proving Ground, MD 21005-5066

Commander  
Chemical Research and Development Center  
U.S. Army Armament, Munitions  
and Chemical Command  
ATTN: SMCCR-SPS-IL  
Aberdeen Proving Ground, MD 21010-5423

Commander  
Chemical Research and Development Center  
U.S. Army Armament, Munitions  
and Chemical Command  
ATTN: SMCCR-RSP-A  
Aberdeen Proving Ground, MD 21010-5423

Director  
Ballistic Research Laboratory  
ATTN: AMXBR-OD-ST  
Aberdeen Proving Ground, MD 21005-5066



Chief  
Benet Weapons Laboratory, CCAC  
Armament Research and Development Center  
U.S. Army Armament, Munitions  
and Chemical Command  
ATTN: SMCAR-CCB-TL (2)  
Watervliet, NY 12189-5000

Commander  
U.S. Army Armament, Munitions  
and Chemical Command  
ATTN: SMCAR-ESP-L  
Rock Island, IL 61299-6000

Director  
U.S. Army TRADOC Systems  
Analysis Activity  
ATTN: ATAA-SL  
White Sands Missile Range, NM 88002

Director  
Defense Advanced Research  
Projects Agency  
ATTN: DARPA-TTO, H. D. Fair, Jr. (5)  
1400 Wilson Boulevard  
Arlington, VA 22209

Commander  
U.S. Army Strategic Defense Command  
ATTN: DASD-DP, LTC N. Cooney (2)  
P.O. Box 15280  
Arlington, VA 22215-0151

System Planning Corporation  
ATTN: Donald E. Shaw  
1500 Wilson Boulevard  
Arlington, VA 22209-2454

Center for Electromechanics  
ATTN: W. Weldon (2)  
10100 Burnet Road  
Austin, TX 78758

Commander  
HQ, AFSC/DLWA  
ATTN: CPT R. Reynolds  
Andrews Air Force Base, MD 20334

Commander  
Naval Sea Systems Command  
ATTN: LCDR Joseph R. Costa  
Washington, DC 20362

Commander  
Air Force Armament Technology Laboratory  
ATTN: DLDG, Timothy Aden  
Eglin Air Force Base, FL 32542-5270

Commander  
U.S. Army Research Office  
ATTN: AMXRO-PR, H. T. Throckmorton  
P.O. Box 12211  
Research Triangle Park, NC 27709

Commander  
Strategic Defense Initiative Organization  
ATTN: SDIO/KEW, BG M. O'Neil (2)  
Washington, DC 20301-7100

Electromagnetic Launch Research, Inc.  
ATTN: Henry Kolm  
Peter P. Mongeau  
625 Putnam Avenue  
Cambridge, MA 92139

Westinghouse Electric Company  
ATTN: Lan R. McNab, Manager,  
EML Engineering  
Marine Division, MS ED-5  
401 E. Hendy Avenue  
P.O. Box 499  
Sunnyvale, CA 94088

Brookhaven National Laboratory  
ATTN: James R. Powell, Jr., Head,  
Fusion Technology Group  
Building 820M  
Upton, NY 11973

General Electrical Company  
ATTN: Alan Wait, Manager,  
EML Systems Technology  
RD-3 Plains Road  
Ballston Spa, NY 12020

END  
FILMED

5-86

DTIC